

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**



(72) CHAMBERS, John George, GB  
(72) TURNER, Graham Andrew, GB  
(71) UNILEVER PLC, GB  
(51) Int.Cl.<sup>6</sup> A61K 7/48, A61K 7/40  
(30) 1996/02/02 (9602111.8) GB  
(54) **COMPOSITION D'HYGIENE CORPORELLE**  
(54) **PERSONAL CARE COMPOSITION**

$$0.5 < \frac{V}{a_o I_c} \leq 1.0; \quad (I)$$

(57) Cette invention se rapporte à une composition d'hygiène corporelle qui se présente sous la forme d'un liquide aqueux comportant (i) une composition liquide contenant deux composants D et E, D étant une molécule ayant une longue chaîne hydrocarbure et un groupe de tête hydrophile et E étant une matière comportant au moins un composé sélectionné dans le groupe constitué par le 3.beta.-stérol, le squalane, le squalène, des saponines ou sapogénines du type stéroïde ou triterpénoïde végétal, les di et tri terpènes du type phytol, rétinol et amyryne, et des mélanges de ces composés, D et E étant présents en quantités représentant respectivement de 0,1 à 10 % en poids et de 0,2 à 12 % en poids de la composition totale, (ii) un tensioactif sélectionné parmi des tensioactifs anioniques, non ioniques, cationiques, zwitterioniques ou amphotères, du savon ou des mélanges de ces composés, et (iii) un auxiliaire de déposition. Ladite composition ne contient pratiquement pas de molécule ayant au moins deux chaînes hydrocarbures et un groupe de tête polaire satisfaisant la relation (I) dans laquelle V est le volume des chaînes hydrocarbures,  $I_c$  est la longueur critique de ces chaînes hydrocarbures et  $a_o$  est l'aire optimale du groupe de tête polaire. Cette composition peut se présenter sous la forme, par exemple, d'un gel pour la douche ou d'un démaquillant pour le visage que l'on applique temporairement sur la peau avant de le retirer soit en essuyant, soit en rinçant.

(57) A personal care composition in the form of an aqueous liquid comprises: (i) a lipid composition comprising two components D and E, where D is a molecule having one long hydrocarbon chain and a hydrophilic head group and E is a material which comprises at least one of a compound selected from 3.beta.-sterol; squalane; squalene; saponins or sapogenins of the plant steroid or triterpenoid type; di and tri terpenes such as phytol, retinol and amyryne; and mixtures thereof, wherein D and E are respectively present at levels within the range 0.1 to 10 wt.% and 0.2 to 12 wt.% based on the total composition; (ii) a surface active agent selected from anionic, nonionic, cationic, zwitterionic, amphoteric surface active agents, soap and mixtures thereof; and (iii) a deposition aid; and the composition is substantially free of a molecule having at least two hydrocarbon chains and a polar head group which satisfies the relationship (I) where V is the volume of the hydrocarbon chains;  $I_c$  is the critical length of the hydrocarbon chains; and  $a_o$  is the optimum area of the polar head group. The composition may be in the form of for example a shower gel or facial cleanser which is temporarily applied to the skin before being removed such as wiping or rinsing it from the skin.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 7/48</b>	<b>A1</b>	(11) International Publication Number: <b>WO 97/27835</b> (43) International Publication Date: <b>7 August 1997 (07.08.97)</b>
(21) International Application Number: <b>PCT/EP97/00199</b> (22) International Filing Date: <b>13 January 1997 (13.01.97)</b> (30) Priority Data: <b>9602111.8</b> <b>2 February 1996 (02.02.96)</b> <b>GB</b> (71) Applicant (for AU BB CA GB IE IL KE LC LK LS MN MW NZ SD SG SZ TT UG only): <b>UNILEVER PLC [GB/GB]; Unilever House, Blackfriars, London EC4P 4BQ (GB).</b> (71) Applicant (for all designated States except AU BB CA GB IE IL KE LC LK LS MN MW NZ SD SG SZ TT UG): <b>UNILEVER N.V. [NL/NL]; Weena 455, NL-3013 AL Rotterdam (NL).</b> (72) Inventors: <b>CHAMBERS, John, George; 11 Central Avenue, Bromborough, Wirral, Merseyside L62 2BS (GB). TURNER, Graham, Andrew; 6 Greenfields Avenue, Bromborough, Wirral, Merseyside L62 6DD (GB).</b> (74) Agent: <b>ELLIOTT, Peter, William; Unilever plc, Patent Division, Colworth House, Sharnbrook, Bedford MK44 1LQ (GB).</b>		(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: <b>PERSONAL CARE COMPOSITION</b>  $0.5 < \frac{V}{a_o I_c} \leq 1.0; \quad (I)$ (57) Abstract  A personal care composition in the form of an aqueous liquid comprises: (i) a lipid composition comprising two components D and E, where D is a molecule having one long hydrocarbon chain and a hydrophilic head group and E is a material which comprises at least one of a compound selected from 3 $\beta$ -sterol; squalane; squalene; saponins or sapogenins of the plant steroid or triterpenoid type; di and tri terpenes such as phytol, retinol and amylin; and mixtures thereof, wherein D and E are respectively present at levels within the range 0.1 to 10 wt.% and 0.2 to 12 wt.% based on the total composition; (ii) a surface active agent selected from anionic, nonionic, cationic, zwitterionic, amphoteric surface active agents, soap and mixtures thereof; and (iii) a deposition aid; and the composition is substantially free of a molecule having at least two hydrocarbon chains and a polar head group which satisfies the relationship (I) where V is the volume of the hydrocarbon chains; I <sub>c</sub> is the critical length of the hydrocarbon chains; and a <sub>o</sub> is the optimum area of the polar head group. The composition may be in the form of for example a shower gel or facial cleanser which is temporarily applied to the skin before being removed such as wiping or rinsing it from the skin.		

- 1 -

PERSONAL CARE COMPOSITION

The invention relates to a personal care composition in liquid or gel form suitable for personal washing. In particular, it is concerned with compositions, intended for washing and/or cleansing of the human skin, which prevent or ameliorate skin dryness, skin wrinkling, chapping and ageing.

It is generally understood that ceramides present within the intercellular lipid lamellae of the stratum corneum play an important role in the production and maintenance of the water permeability barrier of the skin. Ceramides, or substances closely related to them, have been disclosed as components of skin care compositions. In particular, Kao Corporation in GB 2178312 and GB 2213723 discloses the use of natural ceramides extracted from skin in products for topical application. Also, Kao Corporation in EP 227994 and EP 282816 discloses the use of synthetic ceramides, which are similar to their natural counterparts found in skin.

It is also known that, in addition to ceramides, the lipid lamellae comprises sterol and fatty acids (N Y Schurer, P M Elias (1991) Adv Lip Res 24 27-56 Acad Press).

It is believed that one of the causes of skin drying and ageing is a reduction in the amount of lipid contained within intercellular lipid lamella of the stratum corneum. It has been shown in Intermolecular and Surface Forces, (1985) Jacob N Isrealachvili, ed Acad Press, Chapter 16 entitled "Aggregation of Amphiphilic Molecules into Micelles, Bilayers, Vesicles and Biological Membranes" that suitable lipids for forming a bilayer are those having a polar head group and at least two hydrocarbon chains, such that there exists a clearly defined relationship between the volume

- 2 -

occupied by the hydrocarbon chains and the optimum area occupied by the polar head group. This relationship is that:

$$0.5 < \frac{V}{a_o I_c} \leq 1.0; \quad (I)$$

5 where

V is the volume of the hydrocarbon chains

$I_c$  is the critical length of the hydrocarbon chains

$a_o$  is the optimum area of the polar head group.

10 EP-A-556 957 discloses compositions comprising (a) a molecule having at least two hydrocarbon chains and a polar head group which satisfies the relationship (I) above such as ceramides, (b) long chain fatty acids and a third component (c), for  
15 example squalene, which is capable of assisting and stabilising lipid bilayers formed in the composition and where the ratio of a:b:c is from 1:1.5 to 6.0:1.1 to 8.0 respectively. The disclosed compositions are intended for topical application to the human, hair, skin or nails. In  
20 particular they are "leave-on" products which are intended to be permanently applied to the hair, skin or nails.

WO 94/00127 also discloses three component lipid compositions which may be combined with an aqueous phase containing a surface active agent. Like EP-A-557 957, the compositions  
25 disclosed in this reference are intended as "leave-on" products.

The aforementioned prior art compositions are intended as "leave-on" products which are not wiped or rinsed off. A  
30 disadvantage with such compositions is that they may leave the skin feeling oily.

There is a continued need for products which are able to successfully replace depleted lipids and alternative  
35 compositions such as cleansing compositions which are only temporarily applied to the skin before being removed. In our

- 3 -

5 copending patent application PCT/EP95/03967 personal care compositions comprising a lipid composition comprising molecule A) which satisfies relationship (I) above such as a sucrose ester; B) long chain fatty acids; and a third  
10 component C) such as cholesterol which is capable of assisting and stabilising lipid bilayers formed in the composition can be formulated with surfactants to form a personal care composition which is intended to be temporarily applied to the skin before being rinsed or wiped off are disclosed. These personal care compositions provide an effective control of water loss and/or repair of damage to the water barrier layer in the stratum corneum.

15 We have now found that it is possible to formulate compositions in the absence of the aforementioned component A whilst still maintaining the beneficial properties of the prior art compositions i.e. effective control of water loss and/or repair of damage to the water barrier layer in the stratum corneum.

20 Thus, according to the invention there is provided a personal care composition in the form of an aqueous liquid comprising

- 25 i) a lipid composition comprising two components D and E, where D is a molecule having one long hydrocarbon chain and a hydrophillic head group and E is a material which comprises at least one of a compound selected from 3 $\beta$ -sterol; squalene; squalane; saponins or sapogenins of the plant steroid or triterpenoid type; di and tri terpenes such as phytol, retinol and amyrrin; and mixtures thereof, wherein D and E are respectively  
30 present at levels within the range 0.1 to 10wt% and 0.2 to 12wt% based on the total composition;
- 35 ii) a surface active agent selected from anionic, nonionic, cationic, zwitterionic, amphoteric surface active agents, soap and mixtures thereof; and

- 4 -

iii) a deposition aid; and

the composition is substantially free of a molecule having at least two hydrocarbon chains and a polar head group which satisfies the relationship

$$0.5 < \frac{V}{a_o I_c} \leq 1.0; \quad (I)$$

where

V is the volume of the hydrocarbon chains

$I_c$  is the critical length of the hydrocarbon chains and

$a_o$  is the optimum area of the polar head group.

Furthermore, the invention also provides a personal care composition in the form of an aqueous liquid comprising

i) a lipid composition comprising two components D and E, where D is a molecule having one long hydrocarbon chain and a hydrophilic head group and E is a material which comprises at least one of a compound selected from  $3\beta$ -sterol; squalene; squalane; saponins or sapogenins of the plant steroid or triterpenoid type; di and tri terpenes such as phytol, retinol and amyrrin; and mixtures thereof, wherein D and E are respectively present at levels within the range 0.1 to 10wt% and 0.2 to 12wt% based on the total composition;

ii) a surface active agent selected from anionic, nonionic, cationic, zwitterionic, amphoteric surface active agents, soap and mixtures thereof; and

iii) a deposition aid; and

is substantially free of ceramides; pseudoceramides;

phospholipids; glycolipids having a structure of two or more acyl or alkyl long chains suitably containing from 14 to 50 carbon atoms each, attached to a polar group; specific esters of polyethylene glycol; polyglycerol-n-x oleate (CAS 9007-48-1); sorbitan dioleate (CAS 29116-98-1); sorbitan sesquioleate

WO 97/27835

PCT/EP97/00199

- 5 -

(CAS 8007-43-0); long chain alkyl ether versions of phospholipids and glycolipids.

By substantially free is meant that this component, if present, in the compositions of the invention is only present in trace amounts introduced by way of impurities in the other constituents of the composition.

#### Component D

Component D is a molecule having one long hydrocarbon chain and a hydrophilic head group. Preferred materials are straight-chained mono carboxylic acids having 8 to 24 carbon atoms, straight-chained fatty alcohols having 8 to 24 carbon atoms, sugar esters, alkylated sugars and mixtures thereof.

Most preferably the materials are selected from straight chained substantially saturated (i.e. at least 85% saturated) mono-carboxylic fatty acids having 8 to 20 carbon atoms and straight chained saturated fatty alcohols having 8 to 20 carbon atoms. Mixtures of these fatty acids or fatty alcohols may also be used.

Particularly preferred materials are stearic acid and cetostearyl alcohol.

Preferably component D is present at a level within the range 0.2 to 7wt% based on the total composition.

#### Component E

As stated above, E is a material which comprises at least one of a compound selected from  $3\beta$ -sterol; squalane; squalene; saponins or sapogenins of the plant steroid or triterpenoid type; di and tri terpenes such as phytol, retinol and amyirin; and mixtures thereof. Potential sources



- 6 -

of such material consist of wool wax alcohols (lanolin alcohols) and other alcoholic fractions of lanolin.


5 Preferably component E comprises a  $3\beta$ -sterol having a tail on the 17 position and having no polar groups, for example cholesterol, sitosterol, stigmasterol and ergosterol. Commercially available sources of cholesterol include Super Hartolan ex Croda which comprises at least 30% cholesterol.

10 Cholesterol is a vital component of the natural skin lipids constituting the moisture barrier in the stratum corneum.

15 Preferably component E is present at a level within the range 0.3 to 10wt% based on the total composition.


The surface active agent can be selected from any known surfactant suitable for topical application to the human body. Mild surfactants, i.e. surfactants which do not damage the stratum corneum, the outer layer of skin, are particularly preferred.

20 One preferred anionic detergent is fatty acyl isethionate of formula:



25 where R is an alkyl or alkenyl group of 7 to 21 carbon atoms and M is a solubilising cation such as sodium, potassium, ammonium or substituted ammonium. Preferably at least three quarters of the RCO groups have 12 to 18 carbon atoms and may be derived from coconut, palm or a coconut/palm blend.

30 Another preferred anionic detergent is alkyl ether sulphate of formula:



35

- 7 -

where R is an alkyl group of 8 to 22 carbon atoms,  
n ranges from 0.5 to 10 especially 1.5 to 8, and  
M is a solubilising cation as before or  $Mg^{2+}$ .

- 5 Other possible anionic detergents include alkyl glyceryl ether sulphate, sulphosuccinates, taurates, sarcosinates, sulphoacetates, alkyl phosphate, alkyl phosphate esters and acyl lactylate, alkyl glutamates, and mixtures thereof.
- 10 If the surface active agent comprises soap, the soap is preferably derived from materials with a  $C_8$  to  $C_{22}$  substantially saturated carbon chain and, preferably, is a potassium soap with a  $C_{12}$  to  $C_{18}$  carbon chain.
- 15 Sulphosuccinates may be monoalkyl sulphosuccinates having the formula:  $R^5O_2CCH_2CH(SO_3M)CO_2M$ ; and amido-MEA sulphosuccinates of the formula:  $R^5CONHCH_2CH_2O_2CCH_2CH(SO_3M)CO_2M$ ; wherein  $R^5$  ranges from  $C_8$ - $C_{20}$  alkyl, preferably  $C_{12}$ - $C_{15}$  alkyl and M is a solubilising cation.
- 20 Sarcosinates are generally indicated by the formula:  $R^5CON(CH_3)CH_2CO_2M$ , wherein R ranges from  $C_8$ - $C_{20}$  alkyl, preferably  $C_{12}$ - $C_{15}$  alkyl and M is a solubilising cation.
- 25 Taurates are generally identified by the formula:  $R^5CONR^6CH_2CH_2SO_3M$ , wherein  $R^5$  ranges from  $C_8$ - $C_{20}$  alkyl, preferably  $C_{12}$ - $C_{15}$  alkyl,  $R^6$  ranges from  $C_1$ - $C_4$  alkyl, and M is a solubilising cation.
- 30 Harsh surfactants such as primary alkane sulphonate or alkyl benzene sulphonate will generally be avoided.
- 35 Suitable nonionic surface active agents include alkyl polysaccharides, lactobionamides, ethyleneglycol esters, glycerol monoethers, polyhydroxyamides (glucamide), primary

WO 97/27835

PCT/EP97/00199


- 8 -

and secondary alcohol ethoxylates, especially the C<sub>8-20</sub> aliphatic alcohols ethoxylated with an average of from 1 to 20 moles of ethylene oxide per mole of alcohol.

5 Mixtures of any of the foregoing surface active agents may also be used.

10 The surface active agent is preferably present at a total level of from 1 to 45 wt%, preferably 3 to 30 wt%, and most preferably 5 to 20 wt%.

15 It is also preferable that the composition includes from 0.5 to 15 wt% of a cosurfactant agent with skin-mildness benefits. Suitable materials are zwitterionic detergents which have an alkyl or alkenyl group of 7 to 18 carbon atoms and comply with an overall structural formula



25 where R<sup>1</sup> is alkyl or alkenyl of 7 to 18 carbon atoms  
R<sup>2</sup> and R<sup>3</sup> are each independently alkyl, hydroxyalkyl or carboxyalkyl of 1 to 3 carbon atoms

m is 2 to 4


30 n is 0 or 1

X is alkylene of 1 to 3 carbon atoms optionally substituted with hydroxyl, and


Y is -CO<sub>2</sub><sup>-</sup> or -SO<sub>3</sub><sup>-</sup>

35 Zwitterionic detergents within the above general formula include simple betaines of formula:

- 9 -




and amido betaines of formula:




where m is 2 or 3.

In both formulae  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  are as defined previously.  $\text{R}^1$  may, in particular, be a mixture of  $\text{C}_{12}$  and  $\text{C}_{14}$  alkyl groups derived from coconut so that at least half, preferably at least three quarters of the groups  $\text{R}^1$  have 10 to 14 carbon atoms.  $\text{R}^2$  and  $\text{R}^3$  are preferably methyl.


A further possibility is a sulphobetaine of formula:



or



where m is 2 or 3, or variants of these in which  $-(\text{CH}_2)_3\text{SO}_3^-$  is replaced by




$\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^3$  in these formulae are as defined previously.

- 10 -

Preferably the deposition aid is a cationic polymer present in an amount of at least 0.05% by weight of the total composition. It may well not exceed 3% or even 2% of the composition.

5 Various cationic polymers may be used. Examples of cationic polymers include the cationic cellulose ethers described in US Patents Nos. 3816616 and 4272515 and which are available commercially from Union Carbide Corp. under the trade mark  
10 POLYMER JR. Other suitable materials are the cationic polygalactomannan gum derivatives described in US Patent No. 4298494 which are commercially available under the trade mark JAGUAR. An example of a suitable material is the hydroxypropyltrimethylammonium derivative of guar gum of the  
15 formula:



20 where G represents guar gum. Such a material is available under the name JAGUAR C-13-S. This material also has the CTFA designation Guar Hydroxypropyltrimonium Chloride. In JAGUAR C-13-S the degree of substitution of the cationic  
25 groups is about 0.13. Another possible material is that known as JAGUAR C-17 which is similar to JAGUAR C-13-S but has a higher degree of substitution of cationic groups of about 0.25-0.31. A further example of a guar derivative is the hydroxypropylated cationic guar derivative known as  
30 JAGUAR C-16 which as well as containing the above cationic quaternary ammonium groups also contain hydroxypropyl (-CH<sub>2</sub>CH(OH)CH<sub>3</sub>) substituent groups. In JAGUAR C-16 the degree of substitution of the cationic groups is 0.11-0.16 and the moles of substitution of hydroxypropyl groups is 0.8-  
35 1.1.

- 11 -

Other cationic polymers include cationic polyamides such as the low molecular weight adipic acid/diethylene-triamine polyamide and the copolymers of vinylpyrrolidone and dimethylaminoethyl methacrylate quaternised with dimethyl sulphate (Gafquat 755, GAF Corporation) described in US Patent No. 4080310; the graft cationic copolymer containing N-vinylpyrrolidone, dimethylaminoethyl methacrylate and polyethylene glycol described in US Patent No. 4048301; the mineral acid salts of the amino-alkyl esters of homo- and copolymers of unsaturated carboxylic acids having from 3 to 5 carbon atoms described in US Patent No. 4009256; the polymers of etherified starch described in US Patent No. 3186911; and cationic polyacrylamides of the type described in International Patent Application WO95/22311.

The high molecular weight cationic polymers are sold under the trade mark MERQUAT by Merck & Co., Inc. Representative are Merquat 100, a highly charged cationic dimethyldiallylammonium chloride homopolymer, and Merquat (TM) 550, a highly charged cationic copolymer prepared with dimethyldiallylammonium chloride and acrylamide. These materials are designated in the CTFA dictionary as Quaternium-40 and Quaternium-41, respectively.

Preferred cationic polymers include cationic cellulose ethers, cationic polygalactomannan gums and cationic polyacrylamides.

Other typical optional components which may be included in the compositions of the invention are humectants such as glycerol up to 10wt%; opacifiers, preferably at a level of 0.2 to 2.0 wt%; preservatives, preferably 0.2 to 2.0wt% and perfumes, preferably 0.5 to 3.0wt%; bactericides; colourants; antioxidants; skin-feel modifiers; and thickeners and structurants such as swelling clays, cross-linked

- 12 -

polyacrylates, for example, Carbopol (TM) (polymers available from Goodrich) and polyethoxypropylene glycoldioleate.

5 Compositions of the invention may be formulated as products for washing the skin, for example bath or shower gels; hand-washing compositions; facial washing liquids; and skin cleansers.

10 Without being bound by theory it is believed that the deposition aid assists deposition of the lipid composition onto the skin.

15 The composition of the invention is intended, during use, to reduce the permeability of the skin to water, particularly when the skin is dry or damaged, in order to reduce moisture loss and generally enhance the quality and flexibility of the skin.

20 In use, the composition is temporarily applied to the skin directly or by the use of an applicator. It is then removed from the skin such as by rinsing with water, particularly in the case of shower gels, or by wiping off such as with a tissue.

25 The composition according to the invention is preferably formulated as a liquid or gel having a viscosity in the range 1000 to 100 000 mPas measured at a shear rate of  $10\text{s}^{-1}$  and  $25^{\circ}\text{C}$  in a Haake Rotoviscometer RV20.

30 Compositions according to the invention may be prepared by melting the components of the lipid composition. Water is then added to the resultant lipid melt. The surface active agent is melted and then added to the lipid/water mixture. The resultant mixture is cooled before cationic polymer and  
35 optional components are added.

- 13 -

The composition may be packaged in a suitable container from which it can be dispensed directly onto the skin or via an applicator. The invention accordingly provides a closed container containing the personal care composition as herein defined.

The invention also provides for the use in an aqueous liquid personal care composition which is temporarily applied to the skin of

- i) a lipid composition comprising two components D and E, where D is a molecule having one long hydrocarbon chain and a hydrophillic head group and E is a material which comprises at least one of a compound selected from 3 $\beta$ -sterol; squalane; squalene; saponins or sapogenins of the plant steroid or triterpenoid type; di and tri terpenes such as phytol, retinol and amyrrin; and mixtures thereof, wherein D and E are respectively present at levels within the range 0.1 to 10wt% and 0.2 to 12wt% based on the total composition;
- ii) a surface active agent selected from anionic, nonionic, cationic, zwitterionic, amphoteric surface active agents, soap and mixtures thereof; and
- iii) a deposition aid; and

the composition is substantially free of a molecule having at least two hydrocarbon chains and a polar head group which satisfies the relationship

$$0.5 < \frac{V}{a_o I_c} \leq 1.0; \quad (I)$$

where

V is the volume of the hydrocarbon chains  
I<sub>c</sub> is the critical length of the hydrocarbon chains and  
a<sub>o</sub> is the optimum area of the polar head group.



- 14 -

The invention will now be further illustrated by reference to the non-limiting examples.

### Examples

5

In the examples:-

Lauric acid was Prifrac 2922 ex Unichema.

Myristic acid was Prifrac 2940.

10 Ethylene glycol monostearate was Empilan EGMS ex Albright & Wilson.

PEG 6000 distearate was Crothix ex Croda.

15 Cationic polymer was guar hydroxypropyl trimonium chloride was Jaguar C-13-S ex Meyhall except in example 2 where it was Jaguar C-14-S ex Meyhall.

Lanolin alcohol was Super Hartolan ex Croda containing at least 30% cholesterol.

Coco amidopropyl betaine (CAPB) was Tegobetaine C ex Goldschmidt.

20 Sodium lauryl ether sulphate (SLES) was Elfan NS 243S ex Akzo.

Stearic acid was Pristerine 4911.

25 Compositions according to the examples were prepared by melting the components of the lipid composition by heating to approximately 70°C. The lipid components were heated in glycerol when present in the formulations. Water was then added at approximately 70°C. In a separate vessel the surface active agents were heated to approximately 70°C

30 before being added sequentially to the molten lipid mixture. The resultant surface active agent/lipid mixture was mixed and mixing continued whilst it was cooled to room temperature. The cationic polymer, predispersed in perfume, was then added followed by minor components. Finally the pH of

WO 97/27835

PCT/EP97/00199

- 15 -

the composition was adjusted. This method may need minor modifications depending on the precise formulation.

Compositions formulated according to the invention and comparative compositions were tested by visual assessment of skin dryness and erythema of treated skin, measurement of the reduction in water flux (TEWL) and measurement of the skin by a Corneometer.

The experimental procedure employed was as follows. This was carried out on human volunteers.

Both forearms of a human volunteer were washed with a commercially available toilet soap bar three times a day for 7 days. The forearms were wetted with warm water and the shower gel was dispensed directly onto the arm and then rubbed into a lather for 45 seconds. The forearms were then rinsed with warm water for 15 seconds before being patted dry. They were monitored for skin dryness and erythema on a daily basis.

On days 8-12 the volunteers forearms were washed with a composition formulated according to the invention or comparative formulations. (Formulations are given below). The forearms were wetted with warm water and the shower gel was dispensed directly onto the arm and then rubbed into a lather for 45 seconds. The forearms were then rinsed with warm water for 15 seconds before being patted dry. They were monitored for skin dryness and erythema on a daily basis. On day 13 only three washes were performed.

This test procedure can be used to assess the effectiveness of the compositions according to the invention in alleviating dryness produced during the first 7 days of the test.

WO 97/27835

PCT/EP97/00199

- 16 -

A) Skin dryness and erythema

This was assessed visually and scored as below.

5	<u>Skin Dryness</u>	<u>Score</u>
	No visible dryness	0
	Slightly white, barely perceptible	1
	Moderate whiteness:	2
10	Slight patchy uplifting of skin	3
	Sight uplifting of skin, uniform scaling	4

	<u>Erythema</u>	<u>Score</u>
15	None	0
	Barely perceptible	0.25
	Slight	0.50
	Mild/patchy	1.0

20

B) Reduction in water flux (Trans Epidermal Water loss - TEWL)

TEWL measurements were carried out using a Servomed Evaporimeter EP1 on a fixed site on each of the volunteer's arms (arbitrarily set at 5 inches from the wrist). Measurements were taken

25

30

- i) prior to the start of trial;
- ii) immediately prior to first wash on day 8;
- and
- iii) at least 1 hour after the last wash on day 13.

35

A Teflon probe, containing two sensors, was rested on the surface of the skin. These sensors measured the partial water vapour pressure at two distances above the skin

WO 97/27835

PCT/EP97/00199

- 17 -

surface. The read out from the equipment gave the rate of water evaporation from the skin surface.

C) Corneometer readings

Corneometer readings were taken at the same time as the TEWL readings. The procedure involved resting a conductance probe on the skin surface for a few seconds. The value obtained provided a measure of the hydration state of the outer layer of skin.

D) Amount of lipid deposited onto skin

The following method was used to determine the amount of lipid deposited onto the skin of a human volunteer after it had been washed with the compositions according to the invention was determined.

The volunteers washed their forearms with a shower gel. The procedure involved wetting the arm and also the volunteer's free hand with warm water then using the free hand to lather the arm with 0.5 ml of a formulation for 10 seconds, next rinsing for 10 seconds while rubbing with the free hand and then drying the arm with a single pass with a paper towel.

10 minutes after drying the forearm a glass cylinder was applied to two areas of skin on the forearm. The skin enclosed by the cylinder ( $10\text{cm}^2$ ) was extracted three times with 1ml of ethanol. The amount of cholesterol in the extract was determined using an enzymatic kit from Sigma (352-20). This procedure was repeated for each of the formulations to be tested.

Example 1

The following formulations were prepared and tested.

- 18 -

Table I

<u>% by weight</u>			
<u>Formulation</u>	<u>I</u>	<u>C1</u>	<u>C2</u>
SLES	16.0	16.0	16.0
CAPB	2.0	2.0	2.0
Lanolin alcohol	2.50	-	2.5
Stearic Acid	1.25	-	-
Cationic polymer	0.25	0.25	0.25
Glycerol	5.00	5.00	5.00
Water + preservatives + minors <--- to 100 ----->			

The following results were obtained.

WO 97/27835

PCT/EP97/00199

- 19 -

Table II

<u>FORMULATION</u>	I	C1	Statistical Significance
<b>DRYNESS</b>			
Day 8	2.92	2.77	
Day 13	2.33	1.90	p=<0.05
<b>TEWL/g/m<sup>2</sup>/h</b>			
Day 8	6.80	7.66	
Day 13	8.03	7.15	p=<0.05
<b>HYDRATION (CORNEOMETER)</b>			
Day 8	63.21	61.77	
Day 13	58.23	63.97	p=<0.05
n	21		

n is the number of volunteers who took part in the test.

The results demonstrate the clinical benefit for a composition comprising a binary lipid system in terms of dryness, hydration and water barrier properties of skin treated therewith.

For test D the following results were obtained:-

- 20 -

Formulation	Cholesterol ( $\mu\text{g}/10\text{cm}^2$ )
I	13.54
C1	8.19
C2	11.78

The results demonstrate that when both lipid components are present in the composition there is an increase in the level of lipid deposited compared to that found when the compositions contain only one of the lipid components.

#### Example 2

In this example formulation II, according to the invention, and comparative formulation C3 were each independently compared against a commercially available mild surfactant facial wash product (C4). Twenty panellists were used in this test.

WO 97/27835

PCT/EP97/00199

- 21 -

Table III

Component	II	C3
SLES	12.0	12.0
CAPB	7.5	7.5
Glycerol	5.0	5.0
PEG4000	5.0	5.0
Clay	1.5	1.5
Lanolin alcohol	2.5	2.5
Cetostearyl alcohol	1.25	-
Opacifier	0.4	0.4
Isopropyl palmitate	0.5	0.5
Cationic polymer	0.25	0.25
Perfume + minors + water	<---- to 100 ----->	

The following results were obtained.



- 22 -

Table IV

			Statistical Significance
5	DRYNESS	II	C4
	Day 8	2.45	2.40
	Day 13	1.45	1.85
			p=0.06
	DRYNESS	C3	C4
	Day 8	2.60	2.70
10	Day 13	1.80	1.95
			p=0.49 (NS)

The results demonstrate that a composition containing only one of the lipid components, namely component E, is ineffective compared to a system containing both lipid components D and E.

Example 3

In this example formulations III and IV according to the invention were compared against a system from which lipid component E was absent. (Formulation C5). 8 panellists took part in this test.

- 23 -

Table V

Formulation	III	IV	C5
	wt%		
Lauric Acid	7.69	7.96	7.69
Myristic Acid	7.69	7.69	7.69
Stearic Acid	9.62	9.62	9.62
Ethylene Glycol Monostearate	6.00	6.00	6.00
PEG 6000 Distearate	2.00	2.00	2.00
Propan-1,2-diol	5.00	5.00	5.00
Glycerol	13.00	13.00	13.00
Potassium Hydroxide*	-	-	-
Cationic Guar Gum	0.10	0.10	0.10
Lanolin Alcohols	1.50	3.00	-
Perfume and minors	<----- to 100 ----->		

\* Adjusted to a level to ensure that the fatty acids will not be fully saponified giving a level of free fatty acid of at least 0.2wt%.

The following results were obtained:-

- 24 -

Table VI

Formulation	Cholesterol ( $\mu\text{g}/10\text{cm}^2$ )
III	5.021
IV	5.068
C5	3.621

The results demonstrate that cholesterol is deposited from a fatty acid soap based system personal care composition containing lanolin alcohol.

C3709

- 25 -

CLAIMS

1. A personal washing composition in the form of an aqueous liquid comprising
- 5 i) a lipid composition comprising two components D and E, where D is a molecule having one  $C_{6-24}$  hydrocarbon chain and a hydrophilic head group and E is a material which comprises at least one of a compound selected from  $3\beta$ -sterol; squalane;
- 10 squalene; saponins or sapogenins of the plant steroid or triterpenoid type; di and tri terpenes such as phytol, retinol and amyrrin; and mixtures thereof; wherein D and E are respectively present at levels within the range 0.1 to 10wt% and 0.2 to
- 15 12wt% based on the total composition;
- ii) 1-45 wt% of a surface active agent selected from anionic, nonionic, cationic, zwitterionic, amphoteric surface active agents, soap and mixtures thereof; and
- 20 iii) a deposition aid; and

the composition is substantially free of a molecule having at least two hydrocarbon chains and a polar head group which satisfies the relationship

$$25 \quad 0.5 < \frac{V}{a_0 I_c} \leq 1.0; \quad (I)$$

where

V is the volume of the hydrocarbon chains

$I_c$  is the critical length of the hydrocarbon chains and

30  $a_0$  is the optimum area of the polar head group.

C3709

- 26 -

2. A personal care composition according to claim 1  
wherein component D is selected from straight chained  
substantially saturated mono-carboxylic acids having 8  
to 20 carbon atoms and straight chain substantially  
saturated fatty alcohols having 8 to 20 carbon atoms.
3. A personal care composition according to claim 1  
wherein component E comprises cholesterol.
4. A personal care composition according to claim 1  
wherein the deposition aid is a cationic polymer.
5. A personal care composition according to claim 4  
wherein the cationic polymer is selected from cationic  
cellulose ethers, cationic polygalactomannan gums,  
cationic polyacrylamides and mixtures thereof.
6. Use of a composition according to any preceding claim  
in the topical treatment of dry, aged or damaged skin.
7. Use according to claim 6 in which the personal care  
composition is temporarily applied to skin to be  
treated.
8. Use in an aqueous liquid personal washing composition  
which is temporarily applied to skin of
- i) a lipid composition comprising two components D  
and E, where D is a molecule having one  
C<sub>8-24</sub> hydrocarbon chain and a hydrophillic head  
group and E is a material which comprises at least  
one of a compound selected from 3 $\beta$ -sterol;  
squalane; squalene; saponins or sapogenins of the  
plant steroid or triterpenoid type; di and tri  
terpenes such as phytol, retinol and amyirin; and  
mixtures thereof, wherein D and E are respectively

AMENDED SHEET

C3709

- 27 -

present at levels within the range 0.1 to 10wt%  
and 0.2 to 12wt% based on the total composition;  
ii) 1-45 wt% of a surface active agent selected from  
anionic, nonionic, cationic, zwitterionic,  
5 amphoteric surface active agents, soap and  
mixtures thereof; and  
iii) a deposition aid; and

the composition is substantially free of a molecule  
10 having at least two hydrocarbon chains and a polar head  
group which satisfies the relationship

$$0.5 < \frac{V}{a_0 I_c} \leq 1.0; \quad (I)$$

where

15 V is the volume of the hydrocarbon chains  
I<sub>c</sub> is the critical length of the hydrocarbon chains and  
a<sub>0</sub> is the optimum area of the polar head group.